

Topological Classification of Fermi Surfaces

Y. X. Zhao^{1*} and Z. D. Wang^{1,2†}

*Department of Physics, The University of Hong Kong¹ and
Center of Theoretical and Computational Physics, The University of Hong Kong²*
(Dated: December 3, 2012)

In the same spirit of a classification scenario for Hamiltonian used in the random matrix theory, a kind of topological classification of Fermi surfaces is put forward in terms of topological charges. The topological charge of a Fermi surface depends on its codimension and the class that its Hamiltonian belongs to. It is revealed that there exist six types of topological charges that form two groups with respect to the chiral symmetry, with each group consisting of one original charge and two descendants. Moreover, an intrinsic relationship between the symmetry class index and the codimension number is explicitly disclosed, based on which we are able to realize any type of Fermi surface with a high codimension by reducing its codimension to an experimentally accessible one.

PACS numbers: 03.65.Vf, 71.18.+y

I. INTRODUCTION

For a Fermi gas at zero temperature, a Fermi surface (FS) naturally arises as the boundary separating the occupied and empty states in $(\omega, \mathbf{k})$ -space. As is known, when weak interactions/perturbations and disorders are introduced, some Fermi surfaces still survive though the occupation of states may be shifted dramatically, while others are easily gapped. Such kind of FS stability originates from its topological property of the Green's function or the Feynman propagator for fermionic particles, $G(\omega, \mathbf{k}) = (i\omega - \mathcal{H})^{-1}$, which was first pointed out by Volovik in Ref.[1]. Generally speaking, an FS is robust against weak interactions/perturbations and disorders, if it has a nontrivial topological charge that provides the protection; otherwise it is vulnerable and easy to be gapped. The most general case is that a Hamiltonian is not subject to any symmetry, where the topological charge is formulated by the homotopy group $\pi_p(GL(N, \mathbb{C}))$. This general case was addressed in Refs.[1, 2] and analyzed in the framework of the K theory [3]. Notably, real physical systems have normally certain symmetries, making it necessary and significant to develop a corresponding theory for symmetry preserving cases.

It is known that the symmetry of a quantum system can always have either a unitary representation or an anti-unitary one in the corresponding Hilbert space. The ordinary symmetries, such as rotation, translation, and parity symmetries, are unitary ones. If a system preserves a discrete unitary symmetry, its Hamiltonian commutes with the matrix representing the symmetry; while if it has a continuous symmetry, the Hamiltonian commutes with the matrices representing its generators. A theory for the unitary symmetry was established by involving these matrices in Ref.[1, 2]. However, the unitary symmetries commuting with the Hamiltonian are easily broken by weak interactions/perturbations and disorders.

On the other hand, the anti-unitary symmetries, such

as the time-reversal symmetry (TRS) and charge conjugate or particle-hole symmetry (PHS) can usually be preserved in the presence of weak interactions/perturbations and disorders. Thus we are motivated to develop a unified theory to classify the FSs of systems with the two so-called reality symmetries, namely TRS and PHS, which seems to be fundamentally important.

The paper is organized as follows. Classification of Hamiltonian and general results are introduced in Sec.II A. The two complex cases are discussed in Sec.II B and Sec.II C, respectively. In Sec.II D, the non-chiral real cases classified in Tab.[II] are investigated. We illustrate that the two descendants are induced from the original one mentioned in Sec.II B and discuss physical meanings of the corresponding three topological charges. Sec.II E consists of the chiral real cases indicated in Tab.[II]. Finally, a brief summary is presented in Sec.III.

II. CLASSIFICATION OF FERMI SURFACES AND TOPOLOGICAL CHARGES

A. Introduction of Classification

Let us first to introduce how to classify all sets of Hamiltonian involving the above-mentioned two reality symmetries, as done in the random matrix theory[4, 5]. It turns out that the chiral symmetry (CS) has to be introduced for a complete set of classification. If a unitary symmetry represented by $\mathcal{K}$ can anti-commute with the Hamiltonian $\mathcal{H}$, i.e., $\{\mathcal{K}, \mathcal{H}\} = 0$, it corresponds the CS. In fact, the combined symmetry of TRS and PHS is a kind of chiral symmetry. On the other hand, two chiral symmetries can be combined to commute with $\mathcal{H}$, which makes $\mathcal{H}$ diagonal. Thus it is sufficient to consider only one chiral symmetry. Based on these considerations[6–8], a complete classification of all sets of Hamiltonian can be obtained with respect to the three symmetries. As is known, the TRS and PHS are both anti-unitary, and can

	Non-chiral case					Chiral case				
	A	AI	D	AII	C	AIII	BDI	DIII	CII	CI
T	0	+1	0	-1	0	0	+1	-1	-1	-1
C	0	0	+1	0	-1	0	+1	+1	-1	+1
S	0	0	0	0	0	1	1	1	1	1

TABLE I. T and C denote respectively TRS and PHS, with 0 indicating the absence of TRS(PHS) and ± 1 denoting the sign of TRS(PHS). S denotes CS, with 0 and 1 representing the absence and presence of CS.

be expressed in a unified form

$$\mathcal{H}(\mathbf{k}) = \epsilon_c C \mathcal{H}^T(-\mathbf{k}) C^{-1}, \quad C C^\dagger = 1, \quad C^T = \eta_c C, \quad (1)$$

where $\epsilon_c = \pm 1$ denotes TRS/PHS, and $\eta_c = \pm 1$. As a result, each reality symmetry may take three possible types (even, odd, and absent), and thus there are nine classes. In addition, considering that the chiral symmetry may be preserved or not when both reality symmetries are absent, we can have ten classes, as summarized in Tab.[I], which was first indicated in Refs.[4, 5], while may be easily understood in the present framework [6–9].

For a system with spatial dimension d , the Fermi surface is a connected compact submanifold with dimension $d - p$ if the $(\omega, \mathbf{k})$ -space is compactified. We may define the codimension of the Fermi surface as p . Generally, the topological property of a Fermi surface depends on its codimension and symmetry. After the detailed analysis, we are able to classify all classes in an appropriate form illustrated by Tab.[II], which is one of main results of this work. In each case, as seen in Tab.[II], a kind of topological charge is designated to the Fermi surface. Topological charges 0, $\mathbf{Z}$, $\mathbf{Z}_2$, and $2\mathbf{Z}$ imply that they are trivial, an integer, an integer of $\text{mod } 2$, and an even integer, respectively. As illustrated in the table, the ten classes can be divided into real and complex cases according to whether or not they have either of the reality symmetries. On the other hand, they can also be put into the chiral and non-chiral cases depending on whether or not they have the CS. Actually, from this kind of classification, we can find that there are six types of topological charges, which form two groups in terms of the CS, with each group consisting of an original one ($\mathbf{Z}$) and the two descendants (son and grandson) with $\mathbf{Z}_2$ type. Note that $\mathbf{Z}$ and $2\mathbf{Z}$ are given by the same formula Eq.[3] (Eq.[4]) in the chiral (non-chiral) real case. In the chiral or non-chiral real case, all of the four types of FSs can be realized for any given number of codimension. It is also clear that the complex and real cases have different periodicities: one is of a two-fold periodicity, while the other is of an eight-fold one. Obviously, for the complex case, we can introduce a matrix element $C(p, i)$ to denote the topological charges listed in Tab.[II], where $i = \text{odd}(\text{even})$ number corresponds to the class A (AIII), and p is the number of codimension. In this way, $C(p, i)$ satisfies the relation $C(p, i) = C(p+n, i+n)$ with n an integer

	Non-chiral case		Chiral case					
	Complex case							
	A		AIII					
$p \backslash i$	1		2					
1	Z		0					
2	0		Z					
$\vdots$	$\vdots$		$\vdots$					
	Real case							
	AI	D	AII	C	BDI	DIII	CII	CI
$p \backslash i$	1	3	5	7	2	4	6	8
1	Z	Z ₂	2 Z	0	Z ₂	0	0	0
2	0	Z ₂	0	0	Z	Z ₂	2 Z	0
3	0	Z	Z ₂	2 Z	0	Z ₂	0	0
4	0	0	Z ₂	0	0	Z	Z ₂	2 Z
5	2 Z	0	Z	Z ₂	0	0	Z ₂	0
6	0	0	0	Z ₂	2 Z	0	Z	Z ₂
7	Z ₂	2 Z	0	Z	0	0	0	Z ₂
8	Z ₂	0	0	0	Z ₂	2 Z	0	Z
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$

TABLE II. p is the codimension of a Fermi surface and i is the index of symmetry classes. The ten classes are put in two blocks of rows for the complex and real cases that depend on whether they have the reality symmetry (TRS and/or PHS), while they are divided into two blocks of columns for the chiral and non-chiral cases that correspond to whether they have the CS or not. For each complex case, there exists a two-fold periodicity for the topological numbers; while for each real case, there exists an eight-fold periodicity.

and $i \text{ mod } 2$. For the real case, we can also introduce another matrix element $R(p, i)$, where $i = 1, \dots, 8$ denote the classes AI, BDI, D, DIII, AII, CII, C and CI respectively. Intriguingly, we can also have $R(p, i) = R(p+n, i+n)$ with $i \text{ mod } 8$, which establishes an intrinsic relationship between the symmetry class index and the codimension number. More significantly, based on this relationship, we are able to realize any type of FS with a high codimension by reducing its codimension to an experimentally accessible one, *i.e.*, $p = 1, 2, 3$. Actually, the above relation and the two-fold periodicity as well as the eight-fold one are originated from the Bott periodicity of $GL(N, \mathbf{C})$ (to be discussed in next subsection). In particular, the present eight-fold periodicity is induced by the two reality symmetries enforcing on the two-fold Bott periodicity, and thus is different from the eight-fold Bott periodicity for $O(N)$ and $Sp(N)$ [10]. It is worth to point out that the two reality symmetries are essentially different from the CS, which lies in that each of the reality symmetries relates a $\mathbf{k}$ point to the $-\mathbf{k}$ point in $\mathbf{k}$ -space, while the CS acts on every $\mathbf{k}$. This difference leads to the requirement that a chosen submanifold in $(\omega, \mathbf{k})$ -space has to be centrosymmetric around the origin in order to preserve either of the reality symmetries.

B. Class A

Let us start from the most general case, *i.e.*, the class A, in the absence of any symmetry. The basic idea to classify FSs in terms of the topology may be learnt from this case, which was already discussed in Ref.[3]. We here follow it for a pedagogical purpose. The Green's function can be written as

$$G(\omega, \mathbf{k}) = \frac{1}{i\omega - \mathcal{H}(\mathbf{k})}, \quad (2)$$

which is regarded as an N -dimensional matrix. FSs are defined to be connected with the zero energy. Formulating topological charges in terms of the Green's function has an advantage when handling interacting systems, which is addressed in [3, 11]. The dimensionality of the spacetime is D and that of the space is d . The $(\omega, \mathbf{k})$ -space is compactified. Generally the Fermi surface consists of branches of compact manifolds in $\mathbf{k}$ -space. For a specific branch with dimension $d - p$, a p -dimensional sphere S^p can always be picked up from the transverse dimensions to enclose it in $(\omega, \mathbf{k})$ -space. $G^{-1}(\omega, \mathbf{k})$ is nonsingular for each $(\omega, \mathbf{k})$ restricted on the S^p , in other words, a member of $GL(N, \mathbf{C})$. Then $G^{-1}(\omega, \mathbf{k})$ restricted on the S^p can be regarded as a mapping from S^p to $GL(N, \mathbf{C})$. As all these mappings can be classified by the homotopy group $\pi_p(GL(N, \mathbf{C}))$, $G^{-1}(\omega, \mathbf{k})$ on the S^p is in a certain homotopic class. In the so-called stable regime, $N > \frac{p}{2}$, $\pi_p(GL(N, \mathbf{C}))$ satisfies the Bott periodicity[10],

$$\pi_p(GL(N, \mathbf{C})) = \begin{cases} \mathbf{Z} & p \in \text{odd} \\ 0 & p \in \text{even} \end{cases}.$$

Since physical systems are always in the stable regime, only the mappings on S^p for FSs with the odd codimension $p = 2n + 1$ can be in a nontrivial homotopic class, while ones for the even codimension p are always trivial. In other words, FSs with the even codimension p are always trivial and vulnerable against weak perturbations, while ones with odd codimension can be classified by $\pi_p(GL(N, \mathbf{C})) \sim \mathbf{Z}$ and its stability is topologically protected against any weak perturbations for the non-trivial one. We emphasize that since the class A is not subject to any symmetry, the perturbation can be quiet arbitrary. In addition, the concrete shape of the S^p is irrelevant in principle, and the only requirement is that it is an orientable compact submanifold in $(\omega, \mathbf{k})$ -space with dimension p , which is distinctly different from the cases of reality symmetries. The homotopic number of a mapping for an FS is named as the topological charge of the FS, which can be calculated from the following formula,

$$N_p = C_p \int_{S^p} \text{tr} (G dG^{-1})^p \quad (3)$$

with

$$C_p = -\frac{n!}{(2n+1)!(2\pi i)^{n+1}}.$$

The $\mathbf{Z}$ type of FSs in class A is thus obtained. Once an FS is nontrivially charged by this topological charge, it can survive under any weak perturbations in the absence of any symmetry. But the topologically trivial FSs can be gapped due to the perturbations. Both cases appear in ${}^3\text{He}$ [1]. In ${}^3\text{He-A}$ phase, two Fermi points are topologically protected with charge ± 2 respectively, being a quiet stable phase against weak perturbations. But the planar phase of ${}^3\text{He}$ with a Fermi line cannot have a topological charge because its codimension is even number, and thus the phase is easy to be perturbed to ${}^3\text{He-B}$ phase (fully gapped) or ${}^3\text{He-A}$ phase.

We here wish to pinpoint that $G(\omega, \mathbf{k})|_{S^p}$ can also be used to effectively describe an insulator with the space-time dimension p because of the non-singularity on S^p . In this sense, the present topological charges are similar to some order parameters defined for topological insulator (TI)[8, 11, 12]. The FS appeared there, however, can never be a topologically protected one at the phase transition point of a TI.

C. Class AIII

We now turn to consider the case when the system has a chiral symmetry denoted by $\mathcal{K}$, *i.e.*,

$$\{\mathcal{H}, \mathcal{K}\} = 0.$$

Systems with this symmetry have a crucial property: for any eigenstate $|\alpha\rangle$ of H with the energy E_α , $\mathcal{K}|\alpha\rangle$ is also an eigenstate of H but with the energy $-E_\alpha$. The Hamiltonian associated with this symmetry can always be diagonalized in $\mathbf{k}$ -space on the basis that diagonalizes $\mathcal{K}$ and thus can be written as

$$h(\mathbf{k}) = \begin{pmatrix} & u(\mathbf{k}) \\ u^*(\mathbf{k}) & \end{pmatrix}.$$

On the S^p that is chosen to enclose an FS of dimension $d - p$, $h(\mathbf{k})$ also takes this form, which makes the topological charge defined in Eq(3) always vanishing. However $u(\mathbf{k})$ can be regarded as a mapping from S^{p-1} (set $\omega = 0$) to $GL(N/2, \mathbf{C})$, which may be in a nontrivial homotopic class of $\pi_{p-1}(GL(N/2, \mathbf{C}))$. In this sense, there exist nontrivial FSs for the even number of codimension, *i.e.*, $p = 2n$. The homotopic number may be calculated as

$$\nu_p = C_{p-1} \int_{S^{p-1}} \text{tr} (u(\mathbf{k}) du^{-1}(\mathbf{k}))^{p-1}.$$

As the homotopic number is real, it can also be calculated by $u^*(\mathbf{k})$. Thus the homotopic number can also be

Codimension p				
AI	AII	D	C	
$2\mathbf{Z}$	$4n' + 1$	$8n + 1$	$8n - 1$	$4n' - 1$
$\mathbf{Z}$	$8n + 1$	$8n' + 1$	$4n' - 1$	$8n - 1$

TABLE III. The table shows the integer topological charges in the non-chiral real case. n' is any odd integer, while n is any integer. The pair of AI and AII are exactly opposite to each other because they have TRS with different signs. The pair of D and C are also opposite because they have PHS with different signs.

expressed by Green's function:

$$\nu_p = \frac{C_{p-1}}{2} \int_{S^{p-1}} \text{tr} \left(\mathcal{K} (G d G^{-1})^{p-1} \Big|_{\omega=0} \right). \quad (4)$$

The homotopic number is referred to as the chiral topological charge of the FS.

For this $\mathbf{Z}$ type topological charge in class AIII, if it is nontrivial, the FS is stable against any perturbations that do not break the CS. However the topological protection is not as stable as that induced by Eq.(3), because the FS is gapped if ever the CS is broken. For instance, there exists two Dirac points with the codimension 2 in the honeycomb lattice, where the CS is the sublattice symmetry. Whenever the sublattice symmetry is slightly broken, the system will be fully gapped[13]. On the other hand, it has been realized in the cold-atom experiment[14] that the two Dirac points can survive under quiet strong perturbations when the CS is preserved.

D. Non-chiral Real Cases

1. Integer Case

We now address the topological charges of the real non-chiral case consisting of the class AI, D, AII and C. They all have one of the two reality symmetries (TRS or PHS). Both symmetries relate a $\mathbf{k}$ point to the $-\mathbf{k}$ point. Thus if an arbitrary sphere S^p is chosen in $(\omega, \mathbf{k})$ -space, the Green's function restricted on the S^p may break the reality symmetries. The reality symmetries are preserved on the S^p if and only if the S^p is centrosymmetric, which is different from cases of the class A and AIII. Without loss of generality, it is assumed that there is a standard S^p consisting of points with equal distance to the origin. Because of the reality symmetry in every class, the topological number defined in Eq.(3) cannot survive in any odd codimension. For class AI, it can be nontrivial if and only if $p = 1 \pmod{4}$, and in particular, when $p = 4n' + 1$ with n' odd, the topological charge is an even number; while only when $p = 8n + 1$, the topological charge can be any integer. Other classes in the non-chiral case are similar, as summarized in Tab.[III].

The physical meanings of these topological charges are elaborated as follows. The FSs of the same codimension always appear in pairs with opposite $\mathbf{k}$ coordinates, because both reality symmetries relate $\mathbf{k}$ to $-\mathbf{k}$. There exist two possibilities: *i*) pairs of FSs with codimension p are transverse to the same S^p (with singular points) centered at the origin of $(\omega, \mathbf{k})$ -space; and *ii*) an FS with codimension p resides at the origin in $(\omega, \mathbf{k})$ -space. For the first case, two S^p s (nonsingular) can be chosen to sandwich the S^p that is transverse to pairs of FSs with the codimension p . Here the difference of topological charge defined in Eq.(3) on the two S^p s is just the total topological charge of all FSs sandwiched. If the S^p s belong to a case in Tab.[III], the total topological charge can be classified by $\mathbf{Z}$ or $2\mathbf{Z}$. On the other hand, if they are not in Tab.[III], the total topological charge vanishes. We emphasize that this does not mean that no FS can exist. For example, two stable Fermi points exist with the same distance to the origin for *3He-A*[1], which has the PHS, but they have opposite topological charges ± 2 and thus the total topological charge is zero. For the second case, an S^p can be chosen to enclose the FS centered at the origin. If the S^p belongs to a case in Tab.[III], these FSs can be classified by the topological charge $\mathbf{Z}$ or $2\mathbf{Z}$. However, if the S^p belongs to a trivial case, it is concluded that the FS cannot exist stably at the origin, *i.e.*, the gap is opened at the origin under weak perturbations.

2. The Son Case of $\mathbf{Z}_2$

For an FS with the codimension p one less than that of a $\mathbf{Z}$ -type FS in a non-chiral real case, *e.g.* an FS of codimension 8 above one of codimension 9 in class AI in Tab.[II], the Green's function restricted on the chosen S^p can be classified by a $\mathbf{Z}_2$ -type topological charge. The key idea is that $G(\omega, \mathbf{k})|_{S^p}$ can be continuously extended to $\tilde{G}(\omega, \mathbf{k}; u)$ on the disk D^{p+1} by introducing an auxiliary parameter u (ranging from 0 to 1) with the two requirements: *i*) $\tilde{G}(\omega, \mathbf{k}; 0)|_{S^p} = G(\omega, \mathbf{k})|_{S^p}$, and *ii*) $\tilde{G}(\omega, \mathbf{k}; 1)|_{S^p}$ is a diagonal matrix, such that $\tilde{G}_{\alpha\alpha} = (i\omega - \Delta)^{-1}$ for empty bands and $\tilde{G}_{\beta\beta} = (i\omega + \Delta)^{-1}$ for occupied bands, where Δ is a positive constant[11, 12]. The validity of the extension is based on a fact that $G(\omega, \mathbf{k})$ restricted on S^p is always trivial in the homotopic sense in this case. This topological charge in terms of Green's function is formulated as

$$N_p^{(1)} = C'_p \int_{S^p} \int_0^1 du \text{tr} \left((G d G^{-1})^p G \partial_u G^{-1} \right) \pmod{2} \quad (5)$$

with

$$C'_p = -\frac{2(p/2)!}{p!(2\pi i)^{(p/2)+1}},$$

where “ $\sim$ ” has been dropped for brevity. The topological charge is defined in a similar way to the WZW-term

in quantum field theory[15] and the $\mathbf{Z}_2$ character comes from the fact that two different extensions differ to an even integer, as illustrated in Ref.[11].

At this stage, let us look into the physical implications of the $\mathbf{Z}_2$ topological charge. As mentioned in the previous subsection, the chosen S^p is always centered at the origin in $(\omega, \mathbf{k})$ -space because of the reality symmetries, and there exist two possibilities: *i*) pairs of FSs with the codimension p are transverse to the same S^p centered at the origin of $(\omega, \mathbf{k})$ -space; and *ii*) an FS with codimension p resides at the origin in $(\omega, \mathbf{k})$ -space. For the first, the S^p (singular points on it) being transverse to the FS can be sandwiched by two S^p s (nonsingular). Then a difference of the topological charges defined by Eq.(5) on the two S^p s, *i.e.*, the topological charge in the shell specified by the two S^p s, can be classified by $\mathbf{Z}_2$. If the topological charge in the shell is equal to 1 mod 2, the appearance of FSs in the shell is topologically protected against perturbations that do not break the designated symmetry. But if the charge is equal to 0 mod 2, the FSs are vulnerable and a gap can be opened under weak perturbations. For the second, an S^p can be chosen to enclose the FS and the topological charge with the $\mathbf{Z}_2$ classification can be calculated from Eq.(5), and the topologically nontrivial FS is topologically protected against weak perturbations.

3. The Grandson Case of $\mathbf{Z}_2$

For the second descendant with the codimension p , *e.g.* the case of codimension 7 one step before that of codimension 9 in class AI in Tab.[II], a $\mathbf{Z}_2$ -type topological charge can also be defined on the chosen S^p . To define the $\mathbf{Z}_2$ -type topological charge, the $G(\omega, \mathbf{k})|_{S^p}$ is smoothly extended to a two-dimensional torus T^2 parameterized by the two auxiliary parameters u and v (both ranging from -1 to 1) with the three requirements: *i*) $\tilde{G}(\omega, \mathbf{k}; -1, -1)|_{S^p} = G(\omega, \mathbf{k})|_{S^p}$; and *ii*) $\tilde{G}(\omega, \mathbf{k}; u, v)|_{S^p} = \epsilon_c C \tilde{G}^T(\omega, -\mathbf{k}; -u, -v)|_{S^p} C^{-1}$, referring to Eq.(1); *iii*) $\tilde{G}(\omega, \mathbf{k}; 1, 1)|_{S^p}$ is a diagonal matrix, such that $\tilde{G}_{\alpha\alpha} = (i\omega - \Delta)^{-1}$ for empty bands and $\tilde{G}_{\beta\beta} = (i\omega + \Delta)^{-1}$ for occupied bands. The topological charge is defined as[11, 12]

$$N_p^{(2)} = C_{p+2} \int_{S^p \times T^2} \text{tr} (G dG^{-1})^{p+2} \mod 2, \quad (6)$$

where “ $\sim$ ” has been dropped and the C_{p+2} is defined in Eq.(3). As a topological charge of FSs, its physical meanings are analogous to that of Eq.(5).

E. Chiral Real Cases

For the class AIII in Sec.IIC, it is seen that only FSs with an even number of codimension $p = 2n$ can have

Codimension p					
	CI	DIII	CII	BDI	
$2\mathbf{Z}$	$4n'$	$8n$	$4n' - 2$	$8n - 2$	
$\mathbf{Z}$	$8n$	$4n'$	$8n - 2$	$4n' - 2$	

TABLE IV. The table illustrates the integer topological charges in the chiral real case. n' is any odd integer, while n is any integer.

nontrivial topological charges defined by Eq.(4). When additional reality symmetries are considered, each of the classes BDI, DIII, CII and CI has an eight-fold periodicity, as shown in Tab.[II]. Only two of these four topological charges survive by preserving reality symmetries. One is still $\mathbf{Z}$, while the other becomes $2\mathbf{Z}$, as illustrated in Tab.[IV]. As mentioned before, if a Hamiltonian has the CS and one reality symmetry, it has also the other reality symmetry. Thus all of the four classes have three symmetries. From Tab.[IV], it is observed that the classes CI and DIII have the opposite distributions, stemming from their opposite reality symmetry signs shown in Tab.[I]. The same situation happens to the classes CII and BDI.

Similar to the non-chiral case, a $\mathbf{Z}$ chiral topological charge is associated with two descendants: son and grandson. The $\mathbf{Z}_2$ topological charge of the son is originated from the chiral topological charge defined in Eq.(4) in the same spirit of that Eq.(5) is originated from Eq.(3), which is given by

$$\nu_p^{(1)} = \frac{C'_{p-1}}{2} \int_{S^{p-1}} \int_0^1 du \text{tr} \left(\mathcal{K} (G dG^{-1})^{p-1} G \partial_u G^{-1} \Big|_{\omega=0} \right) \mod 2, \quad (7)$$

where $\mathcal{K}$ is the matrix to represent the chiral symmetry, S^{p-1} is the chosen S^p restricted on $\omega = 0$, and G is extended by an auxiliary parameter ranging from 0 to 1 with the same requirements for introducing Eq.(5). The $\mathbf{Z}_2$ topological charge for the grandson can also be defined in the same spirit of writing out Eq.(6). We extend the $G(\omega, \mathbf{k})|_{S^p}$ to a two-dimensional torus T^2 parameterized by two auxiliary parameters u and v (both ranging from -1 to 1) with the four requirements: the first three requirements are the same as those of the non-chiral counterpart in Eq.(6); while the fourth one is that the chiral symmetry is preserved on T^2 , which also permits that either of reality symmetries is applicable in the second requirement. The $\mathbf{Z}_2$ topological charge is written as

$$\nu_p^{(2)} = \frac{C_{p-1}}{2} \int_{S^{p-1} \times T^2} \text{tr} \left(\mathcal{K} (G dG^{-1})^{p+1} \right) \mod 2, \quad (8)$$

where C_{p-1} is defined in Eq.(3).

Physical implications of the topological charges in the present real chiral case is entirely parallel to those of the real non-chiral case, except that the CS is always required.

III. SUMMARY

FSs with an arbitrary number of codimension in all of the ten symmetry classes have been classified appropriately in terms of topological charges. Two types of fundamental topological charges given by Eq.(3) and Eq.(4) have been applied to the classification of locality, while the classification of the other four topological charges has been implemented in a spherically layered pattern. Remarkably, it has been revealed that the topological stability of FSs with a given codimension p can be exactly predicted once the symmetry type of the Hamiltonian in Tab.[I] is known. More significantly, an intrinsic relationship between the symmetry class index and the codimension number has been disclosed. It has been concluded that every nontrivial topological charge implies certain topological stability protected by the corresponding symmetries.

We thank G. E. Volovik for helpful discussions. This work was supported by the GRF (HKU7058/11P) and the CRF (HKU8/11G) of Hong Kong.

* zhaoyx@hku.hk

† zwang@hku.hk

- [1] G. E. Volovik, *The Universe in a Helium Droplet* (Clarendon, Oxford, 2003).
- [2] G. E. Volovik, arXiv:1111.4627 [hep-ph] (2011).
- [3] P. Hořava, Phys. Rev. Lett **95**, 016405 (2005).
- [4] M. R. Zirnbauer, J. Math. Phys. **37**, 4986 (1996).
- [5] A. Altland and M. R. Zirnbauer, Phys. Rev. B **55**, 1142 (1997).
- [6] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008).
- [7] D. Bernard and A. LeClair, J. Phys. A: Math. Gen. **35**, 2555 (2002).
- [8] S. Ryu, A. P. Schnyder, A. Furusaki and A. W. W. Ludwig, New J. Phys. **12**, 065010 (2010).
- [9] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, AIP Conf. Proc. **1134**, 10 (2008).
- [10] M. Nakahara, *Geometry, Topology, and Physics* (Hilger, London, 1990).
- [11] Z. Wang, X. L. Qi and S. C. Zhang, Phys. Rev. Lett. **105**, 256803 (2010).
- [12] X. L. Qi, T. L. Hughes, and S. C. Zhang, Phys. Rev. B **78**, 195424 (2008).
- [13] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
- [14] L. Tarruell, D. Greif, T. Uehlinger, G. Jotzu and T. Esslinger, Nature **483**, 302 (2012).
- [15] E. Witten, Nucl. Phys. B. **223**, 422 (1983).